

Four-year-contraception rates of mares treated with single injection porcine zona pellucida and GnRH vaccines and an intrauterine device.

Gary Killian^{1,3}, David Thain², Nancy K. Diehl¹, Jack Rhyan³ and Lowell Miller³

¹J. O. Almquist Research Center, The Pennsylvania State University, University Park, PA 16801

²Department of Animal Biotechnology, University of Nevada-Reno, Reno, NV 89557

³ National Wildlife Research Center, USDA-APHIS-WS, Ft. Collins, CO 80521

Corresponding author:

Gary Killian, Ph.D.

USDA-APHIS-WS

7225 Las Vistas Drive

Las Cruces,

NM 88005

Cell 505 636-3151

Email: garykillian@gmail.com

Abstract:

We evaluated the multiyear contraceptive efficacy of the GnRH vaccine GonaCon™, the PZP vaccine SpayVac™, and the human intrauterine device (IUD) 380 Copper “T” in mustang mares provided by the State of Nevada. Eight untreated control mares were compared to 12 mares treated with SpayVac™, 16 mares treated with GonaCon™ and 15 mares treated with the copper-containing IUD. Rates of contraception for years 1, 2, 3 and 4 respectively for SpayVac™ were 100%(12/12), 83%(10/12), 83%(10/12) and 83%(10/12), rates for GonaCon™ were 94% (15 /16), 60% (9/15), 60% (9/15) and 40%(6/15) and rates for IUD-treated mares were 80% (12/15), 29% (4/14),14% (2/14) and 0% (0/14). Antibody titers against PZP and GnRH declined over the four year study. For mares given SpayVac™, uterine oedema was commonly observed.. IUDs were visible by ultrasonography in non-pregnant mustang mares suggesting that pregnant mares did not retain their IUD. IUD retention may be a function of uterine size; pony mares with IUDs had high retention and contraception rates for 4-5 years. We conclude that long-term contraception of mustang mares with a single shot of either the SpayVac™ or GonaCon™ vaccine is possible.

Running Head: Long-term contraception of mustang mares

Key Words: contraception, feral horse, GnRH vaccine, intrauterine device, SpayVac-PZP vaccine

INTRODUCTION

Overpopulation of wild horses is a significant concern in the western United States (Fisher 1983). In Nevada, where most of the wild horses are located, populations grow at a rate of 15-20% a year on State lands, while their range continues to shrink. Current management strategies of removal and adoption are expensive, logistically challenging, and minimally if at all effective in reducing and maintaining wild horse populations at a desired level. Conflicting interests associated with increased movement of people into wild horse ranges, sympathy to maintain wild horse populations because of their historic and cultural importance, competition among horses and indigenous plant and wildlife species, as well as ranching interests are issues impacted by wild horse overpopulation. Controlling fertility of free-ranging horses is considered a viable option for population control. However, this approach has many challenges for which solutions have been elusive. Ideally, methods for contraception of wild horses should be safe and potentially reversible, effective for several years, practical to administer and of reasonable cost and have minimal affect on reproductive or harem behavior. Immunocontraceptive vaccines have garnered considerable attention in recent years as a means to address problems of overabundant wildlife and feral species (Fagerstone et al. 2002, Delves and Roitt 2005, Naz et al. 2005). Two immunocontraceptive vaccines which have been used in a variety of species and for which data exist on their safety and efficacy are porcine zona pellucida (PZP) vaccine (Kirkpatrick et al. 1992, Kirkpatrick et al. 1995, Miller et al. 1999, Miller et al. 2001, Turner and Kirkpatrick 2002, Kirkpatrick and Turner 2002, Kirkpatrick and Turner 2003, Curtis et al. 2007) and gonadotropin releasing hormone (GnRH) vaccine (Miller et al. 2000, Miller et al. 2004, Killian et al. 2006a, Massei et al. this volume). Fertility control of mares using existing porcine zona pellucida (PZP) vaccines has been shown to be safe and effective for up to ten or more years. However, PZP

vaccine preparations used in wild horses have required revaccination every year or two to maintain infertility (Turner et al. 2000, Turner et al., 2002). Limited data exist for the use of GnRH vaccines in mares to control fertility, ovarian function or behavior (Dalin et al. 20002, Killian et al. 2004, Imboden et al. 2006, Killian et al. 2006b, Elhay et al. 2007). Information on multiyear efficacy and effects following a single injection of a GnRH vaccine in mares is lacking. Regardless of the contraceptive vaccine considered, most formulations that have been used do not appear to be effective for the long term without revaccination. Revaccination of mustangs involves considerable expense, manpower, and horse handling to maintain infertility. If a single injection, multiyear contraceptive were available, it may be possible to achieve effective population reduction, and reduce costs and risks associated with frequent horse handling.

We initiated a study in the fall of 2002 and the spring of 2003 to compare the multiyear contraceptive efficacy of a single-shot contraceptive vaccine directed at gonadotropin releasing hormone with that of a single-shot vaccine directed at the zona pellucida of the ovum. We selected GonaConTM to test as the GnRH vaccine based on our positive experiences with it as a single-injection vaccine in deer and other species (Miller et al. 2000, Miller and Killian 2001, Miller et al. 2004, Killian et al. 2006c, Fagerstone et al. this volume). We selected SpayVacTM as the PZP vaccine to test in mares based on our unpublished experiences with it as a single-injection vaccine in white-tailed deer and the reports of multiyear efficacy in harbor seals (Brown et al. 1997) and fallow deer (Fraker et. al. 2002). Both vaccines were administered with AdjuVacTM, an adjuvant developed at the National Wildlife Research Center. In addition, based on preliminary studies with pony mares, we evaluated the use of an intrauterine contraceptive device (IUD), the 380 copper “T” which has been shown to be safe and efficacious in humans for

multiple years (Fortney et al 1999, Wu, et al. 2000). Preliminary results of these studies with Nevada mustangs have been reported earlier (Killian et. al. 2004, 2006b). This paper reports results for these contraceptive approaches after four years of study with Nevada mustangs and 5 years of observation on the use of IUD's in pony mares.

MATERIALS AND METHODS

Animals

Horses for the study were provided by the State of Nevada and were maintained at the Nevada State Penitentiary, located at Carson City. The studies were approved by the Institutional Animal Care and Use Committee of the Pennsylvania State University. A total of 51 mares and three stallions were used. The mares weighed between 225-360 kg and their ages were estimated to be between one and a half to 12 years of age at the time the treatments were given. All mares except the 18 month old filly were observed in the wild to foal normal healthy foals.

For jugular blood sampling and vaccinations, the mares were run into a hydraulic chute and haltered. Vaccines were given intramuscularly in the left lateral neck. Mares were chemically restrained for IUD placement and for pregnancy evaluations by ultrasound or palpation.

Chemical restraint was achieved with an initial intravenous injection of a mixture of 1ml Dormosedan® (detomidine hydrochloride) 10mg, 2 ml xylazine 200 mg, and 2 ml acepromazine 20 mg to produce sedation, followed in 5 min with a second intravenous injection of one bottle of Telazol® (250 mg Tiletamine base, 250 mg Zolazepam base) resuspended in 3 ml xylazine 300mg for anesthesia. This regimen typically gave between 15 and 45 min of anesthesia.

Ten pony mares and two pony stallions belonging to the Department of Dairy and Animal Science of the Pennsylvania State University were used to develop methods and evaluate the use of intra-uterine devices (IUD) as a means to block fertility. The mares ranged in age from 18 months to 12 years when the treatments were administered. Seven of the mare ponies were pastured with a stallion during the entire five year study, except when treatments were being administered or when data were collected. They weighed between 204-306 kg, Three of the mares were on the study for only one year and were pastured with a stallion for two months during the breeding season. They weighed from 281-391 kg.

Ponies were haltered and restrained in a chute for examination and data collection. If sedation was necessary, mares were given intra venous 0.5-1.5 ml xylazine (50-100 mg) and 0.2-0.5 ml butorphanol, or 0.1.-0.25 ml Dormosedan® (10-25 mg) and 0.2-0.5 ml butorphanol (20-50mg).

Treatments

For the Nevada study vaccinations were given in March of 2003. Eleven mares were given a single-shot GnRH vaccine containing 1800 µg of GonaCon™ and four mares were given a single shot GnRH vaccine containing 2800 µg GonaCon™, 12 mares received a single shot PZP vaccine containing 400 µg SpayVac™ and eight mares were assigned to be untreated controls. Copper-containing 380 “T” IUDs were placed in the uterus of 15 mares transcervically, in October of 2002. The SpayVac™ PZP vaccine was provided Dr. Robert Brown (Brown et al., 1997), who developed the vaccine. The GonaCon™ vaccine was provided by the National

Wildlife Research Center, USDA-APHIS-WS. Both SpayVacTM and GonaConTM were made into an emulsion with AdjuVacTM adjuvant (Miller et al., 2004) and injected as a 1 ml dose. To evaluate reproductive capacity treated and control mares were randomly assigned to two breeding groups. “Band Stallions” that had been observed to sire multiple generations in the wild were selected to be pen stallions. Most mares were maintained in the same pen throughout the study, but if behaviour issues resulted in mares fighting, the less dominant mare was moved to the other pen. Mares were penned with a fertile stallion for a breeding trial typically lasting from June through September of each year. If mares failed to breed, or failed to become pregnant or foal during the breeding season, they were considered infertile.

Ponies were used to evaluate three different types of IUDs for ease of placement, retention and efficacy. The 380-copper T and the GyneFix IUD's were purchased from Family Planning Sales Limited, Littlemore, Oxford, UK. The ring IUD was fabricated from Silastic tubing 2.5 mm o.d. which was used to create a ring size of approximately 2.5 cm. Prior to closing the ring with Silastic cement, five or six small copper cable clamps were threaded over the tubing for inclusion in the ring.

In preliminary studies we attempted direct finger insertion of the IUDs into the mare uterus after dilation of the cervix with two fingers. Because this approach was somewhat cumbersome, time consuming and did not easily ensure placement of the IUD deep within the uterine lumen we attempted to use the IUD insertion devices which came with the human IUDs. However, both of these insertion devices were unsatisfactory for the mare uterus. The insertion device for the copper “T” was too short to traverse the mare vagina and cervix. Likewise, the GyneFix IUD the

insertion device, which is intended to attach the IUD to the uterine endometrium with a monofilament, was also too small for the mare reproductive tract. Consequently, we modified disposable large animal uterine swabs to accommodate the IUDs. The modified swabs enable the successful deposit of the IUDs transcervically into the uterine lumen of sedated ponies or anesthetized Nevada mares. The insertion device containing the IUD was placed between the two fingers used to dilate the cervix and guided into the uterine lumen where the IUD was discharged. With minimal practice the time needed to clean the perineum, palpate the cervix by hand, and then insert the device was 2-5 min, with actual insertion of the rod and placement of the device requiring about 30-60 sec.

For the research trial, all IUDs were placed into the uterus transcervically following dilation of the cervix with one or two fingers. As detailed in the results, several animals received more than one type of IUD during the course of the study. If a mare became pregnant after receiving an IUD she was either allowed to go to term, or in the case of two mares where the pregnancy was 60 days or less, the pregnancy was terminated with prostaglandin F₂α. Within 2-3 weeks of pregnancy termination or parturition, mares were treated with another IUD. As a result, the GyneFix IUD was evaluated in 4 mares, the copper ring in 4 mares and the 380 copper “T” in 7 mares. Two untreated mares served as a control for each breeding season.

Observations

Blood samples were collected from treated Nevada mares once or twice a year. However, the only observations consistently made on the eight control mares were general health, body condition and foaling. In mid- to late-October of each year treated mares were examined by

rectal ultrasonography for pregnancy, IUD retention and uterine inflammation. Pregnancy was established by ultrasonography by observance of an embryonic vesicle, a fetus or in the case of later gestation, rectal palpation of a fetus. These observations were later confirmed by birth of a foal. In a few cases where the behaviour of the mare prevented ultrasonography or rectal palpation, pregnancy was determined later by the birth of a foal. General health and body condition, and uterine oedema that may be associated with oestrous cycle changes or presence of an IUD were noted. Uterine oedema in healthy mares is an indication that she is in heat and that she is under the influence of oestrogen produced by ovarian follicles (Sample 1997). All blood samples were assayed for oestradiol, progesterone and antibody titers to the contraceptive vaccines (Miller et al. 2000, 2001). One mare receiving the GonaConTM vaccine and one mare with an IUD died after the first breeding season of causes not related to the treatments.

From April through November of 2002-2005 ponies were gathered and examined by ultrasonography every 4-6 weeks. During the 2004 breeding season, daily observations were also made from April through August on the breeding and harem behavior of a group of ponies consisting of one stallion, two control mares, one mare with a GyneFix IUD, two mares with the ring IUD and four mares with the “T” IUD. For 2006, ponies were pastured for the entire year and the only observations made were for general health status and foaling rates. In April of 2007, ultrasonography was performed to check for IUD placement, pregnancy and any contraindications.

Statistical evaluation

Data from hormone assays were subjected to one way analysis of variance by treatment.

Differences between means were detected by two-sided T test. Mean values are reported plus or minus standard error. Regression analyses were used to evaluate changes in antibody titers during the study.

RESULTS

Nevada mustangs

Foaling data for the eight control mares for years one through four were 75% (6/8 mares), 75% (6/8 mares) 88% (7/8 mares) 100% (8/8 mares), respectively. Rates of contraception for SpayVacTM-treated mares for years one, two, three and four were 100%, 83%, 83% and 83%, respectively (Fig. 1). Average anti-PZP titers for the fall bleed for mares that were contracepted were sustained considerably above the titers of mares that became pregnant (Fig. 2). Although regression analysis indicated a significant decline of anti-PZP titers for the first three years of the study ($p < .05$), average titers tended to increase in the fourth year. Compared to titers in the third year, individual titers increased in eight of the ten contracepted mares, with the remaining two mares having titers similar to the titers they had in the third year. For all years, serum progesterone values (ng/ml) averaged 1.3 ± 0.40 for non-pregnant mares during the October bleed compared to 17.5 the average for two pregnant mares. We were unable to obtain serum oestradiol values for the two SpayVacTM treated mares that became pregnant, but the average serum oestradiol concentrations (ng/ml) for the cycling females at the fall bleed was 25.9 ± 2.9 .

Rates of contraception for GonaConTM treated mares were 93%, 64%, 57% and 43% for years one through four (Fig. 1) For contracepted mares, regression analyses indicated there was

significant decline of average antibody titers over the last three years ($p < .01$). Nevertheless, anti-GnRH titers of non-fertile mares were significantly greater ($p < .01$) than average titers for mares that became pregnant during the study (Fig. 3). Serum progesterone concentrations (ng/ml) for non pregnant GonaConTM treated mares at the fall bleed averaged 0.3 ± 0.1 which was significantly lower ($p < 0.01$) than progesterone concentrations of pregnant mares treated with GonaConTM (7.7 ± 1.5). Average serum oestradiol concentration (pg/ml) for the non-pregnant GonaConTM treated mares for all years was 32.6 ± 6.7 which was significantly less ($p < .001$) than concentrations detected in pregnant GonaConTM treated-mares ($1,179 \pm 225$).

Contraception rates for IUD-treated mares were 80%, 29%, 14% and 0%, respectively in years one through four of the study (Fig. 1). Average serum progesterone for all years at the fall bleed was significantly lower ($p < 0.017$) for non-pregnant IUD-treated mares ($4.3 \pm 1.4 \text{ ng/ml}$) than for pregnant IUD mares ($11.3 \pm 2.7 \text{ ng/ml}$). The average serum oestradiol concentration of $62.5 \pm 34.2 \text{ pg/ml}$ for all years for non-pregnant IUD-treated mares was significantly less ($p < .001$) than the oestradiol serum concentrations of IUD-treated mares that became pregnant ($1,907 \pm 505$).

Comparing fall serum progesterone values among treatments for non-pregnant mares, differences between GonaConTM and SpayVacTM-treated mares were not significant. However, progesterone values in non-pregnant GonaConTM-treated mares and SpayVacTM-treated mares were significantly lower ($p < .02$) than values in non-pregnant IUD-treated mares (Fig. 4). In contrast, serum progesterone values at the fall bleed were lower in pregnant GonaConTM-treated mares than in pregnant-SpayVacTM or pregnant -IUD treated mares (Fig 5). These differences were significant between IUD and GonaConTM-treated mares ($p < .025$), but because only two

SpayVacTM-treated mares became pregnant, statistical comparisons could not be made to the other treatments. There were no significant differences in serum oestradiol concentrations in non-pregnant mares among treatments ($p=.13$) and differences between serum oestradiol concentrations of pregnant IUD-treated and pregnant GonaConTM-treated mares were not significant ($p=.1$).

Observations made by ultrasonography usually enabled visualization of IUD location and the presence of uterine oedema or luminal fluid. In most instances IUDs were not observed in mares that were pregnant. When uterine oedema was observed it was recorded by treatment the mare received (Table 1). These values were compared to the expected incidence of uterine oedema during oestrus based on a normal mare oestrous cycle. We assumed that within a normal 21 day oestrous cycle, 5-7 days of the cycle would be in oestrus, and the remaining 14-16 days the mare would be in dioestrus (Crowell-Davis 2007). Therefore, in a random sample of mares taken from a normal population, approximately 25-30% would be expected to be in oestrus and to have oedematous uteri during the breeding season. Non-pregnant mares treated with GonaConTM had rates of uterine oedema similar to the expected rate of 25-30% for normal non-pregnant cycling mares (Table 1). Likewise, IUD-treated mares in the first year had rates of uterine oedema similar to the expected rate, although in subsequent years too few non-pregnant mares remained in the IUD-treated group to obtain a reliable estimate. In contrast, SpayVacTM-treated mares had high rates of uterine oedema during all four years of the study (Table 1).

Ponies

A summary of the IUDs inserted into the pony mares is provided in Table 2. Only one of the four

mares successfully retained the GyneFix IUD for a five year period. Likewise, the copper-containing Sialistic ring was retained in only one mare for at least a year before she was sold. The copper “T” device provided the greatest rates of contraception and retention with several mares having the device in place for 3-5 years. We were only able to observe two of the mares (Table 2, Wanda, Remy) for one year before they were removed for the study.

Regardless of the type of IUD, the usual reason for failure was pregnancy. In most instances, we assumed the pregnancy resulted from failed retention of the IUD, since we were unable to visualize the IUD by ultrasonography, or to identify it in the afterbirth when observed. In one case, we observed a 60-day pregnancy, but in a subsequent examination six weeks later the mare was not pregnant and the IUD was seen, suggesting the pregnancy was aborted. In one case, a copper “T” was removed because of pyometria. The recovery of that mare was unremarkable.

Two mares (Table 2, Libby, Maddie) were tested with all three IUD types. Both mares had the longest contraception rates with the copper T. It is noteworthy that most mares given the copper “T” IUD remained contracepted for multiple years. However, one mare (Libby) became pregnant within a year, regardless of the IUD type.

Observations made from April through August 2004 on the breeding behavior of three non-pregnant mature mares equipped with a Copper “T” IUD (Dewdrop, Maddie, Godiva) indicated that they had four or five oestrous cycles. The one mature mare equipped with a GyneFix IUD (Sprite) had six oestrous cycles and the one filly equipped with a “T” IUD at 11 months of age had three oestrous cycles. These numbers were within the normal range of oestrous cycles for

pony mares at the Penn State University facility.

Discussion

As with other species, population management of wild horses presents a specific set of challenges for a contraception method to meet in order for it to be of practical use. Assuming the contraception method does not pose serious problems to the health, behavior or well being of the animal, two factors of considerable importance for contraceptive application in wild horses are long-term efficacy and whether the approach is easy to use. Wild horses need to be gathered from their range and managed under safe conditions for them to be hand injected with an immunocontraceptive vaccine. This approach has routinely been used for horses in the western United States. However, it is time consuming, expensive and despite best efforts, not free of risk of injury to the horses or the human handlers. Although remotely darting individuals is possible (Kirkpatrick et al. 1990, Turner et al. 1996), it is not practical for vaccinating large numbers of horses in the western United States. For example, an internal cost-benefit analysis performed by the Nevada Department of Agriculture (D. Thain, unpublished) for the Nevada Virginia Range horse management area, consisting of over 145,000 hectares of mountainous arid rangeland, concluded that equine bands did not aggregate in any one area to enable cost effective darting. Because the Nevada Virginia Range is typical of many horse management areas in the western United States, we believe having a single application contraceptive approach that is effective for multiple years would minimize or remove the need for additional gathers in these areas for revaccination in subsequent years. To date, a single injection vaccine with multiple years of efficacy has not been available for horses despite considerable efforts of investigators working on the problem (Turner et al. 2000, Turner et al. 2002, Liu, et al. 2005). The present study

provides evidence that multiple years of contraceptive efficacy can be achieved with a single shot immunocontraceptive vaccine in the mustang mare. SpayVacTM was shown to have the greatest contraception rate. For the four years of study, only two of the twelve SpayVacTM-treated animals had foals. This rate for a single vaccination far exceeds what has been reported by others for wild horses, although SpayVacTM has been shown to have long term efficacy in other species (Brown et al. 1997, Fraker et. al. 2002). Mare contraception with SpayVacTM was associated with antibody titer, since the two mares which became pregnant had titers much below the average titer of contracepted mares. Although the average titer for the non-pregnant SpayVacTM-treated mares declined in years two and three relative to year one, the titers actually increased in the fourth year in the majority of the contracepted mares. This suggests that self boosting of the immune response may occur (Perry et al., 2006), perhaps as a result of the seasonality of mare reproduction. It is thought that self boosting occurs in the draining lymph node (Burton et al. 1994) as antigen is released from the follicular dendritic cell when the antibody flowing through the draining lymph node drops to a certain level. The released antigen then provides restimulation of antibody production. It is also possible that as the mare returns to breeding condition from a period of anoestrus, new zona pellucida proteins are produced with the initiation of follicle development which restimulate an immune response. If the titer at which mares became pregnant is assumed to be the set point for pregnancy to occur, we would predict that the mares currently contracepted with SpayVacTM will remain infertile for several more years. It is also possible that some of the mares may remain infertile indefinitely.

Compared to GonaConTM and the IUD treatments, mares treated with SpayVacTM had a greater incidence of uterine oedema than would be predicted for normal cycling mares. Uterine oedema

is associated with a predominance of oestradiol (Sample 1997), although there was no significant difference among treatments in oestradiol values for non-pregnant mares. However, expression uterine oedema during the normal oestrous cycle is also influenced by the absence of serum progesterone (Crowell-Davis 2007). Relative to non-pregnant IUD-treated mares which we found to have normal cycle lengths in the pony study, the serum progesterone values were significantly lower in SpayVacTM-treated mares than in IUD-treated mares. Although we were unable to make observations on oestrous cycle length for the Nevada mares in this study, it has been reported that mares treated with PZP vaccine tended to have normal cycle length (Kirkpatrick et al. 1997, Powell 1999) although urinary oestrogen has been reported to be lower (Kirkpatrick et al. 1992), along with a short-term high incidence of persistent corpora lutea (Liu et al. 2005).

Serum oestradiol concentrations determined for SpayVacTM- treated mares suggest that some follicular development occurred. However, for SpayVacTM-treated mares showing some evidence of follicular development based on serum oestradiol, ovarian pathologies may be involved. Ovarian pathologies have been recorded for several species given PZP vaccines (Skinner et al. 1984, Mahi-Brown et al. 1988, Kirkpatrick et al. 1992, Lou et al. 1996, Stoops et al. 2006, Curtis et al. 2007), and are summarized as a disruption of folliculogenesis, depletion of primary oocytes and an infiltration of leucocytes. Given that serum progesterone was significantly lower in SpayVacTM-treated mares than in IUD-treated pony mares known to be having oestrous cycles of normal length, we suggest that SpayVac-treated mares may undergo some follicular development, but fail to ovulate and or develop a normal corpus luteum.

Literature reports of circulating progesterone concentrations for untreated mares indicate considerable variation of values. Although concentrations of serum progesterone of <1 ng/ml are generally associated with oestrus, the concentrations progressively increase to high dioestrous values by days 5-7 and are sustained there until days 13-14. According to Ginther (1992) the range of means for 10 publications assaying serum progesterone during mare dioestrus was 4-22 ng/ml. Clearly, an average exceeding 4 ng/ml for the IUD-treated mares suggests that the majority of IUD mares were in dioestrous when they were sampled. This is what would be expected when randomly sampling a population of mares. In contrast, an average of 1.3 ng/ml for SpayVacTM-treated mares suggests that the majority of those mares were not in dioestrus. Whether they were in oestrus or transitioning into or out of dioestrus cannot be determined with only one sampling point for each mare. Nevertheless, we believe that the published values for average progesterone values during dioestrus, plus the reports of ovarian pathologies associated with PZP vaccines supports the notion that the lower progesterone values in SpayVacTM-treated mares may be due to a failure to form or maintain a normal corpus luteum. It is possible that antibodies to the zona pellucida prevent follicles from developing to normal ovulatory size and formation of the corpus luteum. Aside from these characteristics, there was no other evidence of contraindications associated with the SpayVacTM treatment.

Mares receiving a single vaccination of GonaConTM showed a high degree of contraception the first year, but this rate gradually declined to less than half after 4 years. This decline was associated with a gradual decline in antibody titer to GnRH over the same period. Unlike what was seen with SpayVacTM titers, there was no evidence for a self-boosting effect which occurs when the native protein is produced. The reason for this difference in response to the two

immunogens is unknown, but it may relate to the fact that the PZP immunogen is a large glycoprotein compared to the GnRH decapeptide. Nevertheless, while the contraceptive efficacy of GonaCon™ was not as impressive as SpayVac™, the GonCon™ results exceed rates of contraception reported by others for mares using other single injection contraceptive vaccines (Turner et al. 2000, Turner et al. 2002).

The incidence of uterine oedema for GonaCon™-treated mares was similar to what would be predicted in a population of normal cycling mares. This suggests that these mares may have some degree of oestrous cycle activity, although it is difficult to make firm conclusions. Theoretically, we would predict minimal steroid hormone production in GonaCon™-treated mares, if we assume that gonadotropin releasing hormones were inactivated by antibody to the vaccine. However, plasma oestradiol and progesterone concentrations in GonaCon™-treated mares were similar to those in the other treatment groups. Because we have no direct observations on the reproductive behavior of the GonaCon™-treated mares, we cannot say whether these females expressed oestrus or showed evidence of an oestrous cycle. However, our unpublished data from white-tailed deer treated with GonaCon™ suggests that GonaCon™ does inhibit expression of oestrus at least in the first year or two following vaccination.

The presence of serum oestradiol in GonCon™-treated mares is contrary to the notion that the GnRH is the sole regulator of follicle stimulating hormone (FSH) secretion by the pituitary gland leading to stimulation of follicle development and estrogen secretion. The GonaCon™ used in this study is prepared with the luteinizing hormone releasing hormone (LHRH) peptide (Levy et al. 2004) which has been shown to stimulate both luteinizing hormone (LH) and FSH secretion.

However, there is also evidence in several species for a follicle stimulating hormone releasing hormone (FSHRH) which specifically stimulates FSH secretion by the anterior pituitary gland (McCann et al. 1993, Yu et al. 1997, McCann et al. 1998, Padmanabhan and McNeilly 2001, McNeilly et al. 2003). If an FSHRH exists in the mare, it could explain the presence of serum oestradiol in infertile GonaConTM-treated mares. Antibodies to LHRH in the serum would act to block some follicular development and the LH surge associated with ovulation, but some follicular development and oestrogen production would also occur in response to FSHRH and FSH secretion.

Contraception results for the Nevada mares treated with the IUD were encouraging in the first year of the study, but the performance was poor for the remainder of the study.

When IUDs were visualized, there was no evidence of uterine pathology as assessed by ultrasonography. It has been reported that mares implanted with a Sialistic ring IUDs were infertile for one year, but that device was associated with a uterine inflammatory response (Daels and Hughes 1995). In our study we did not see evidence of a uterine inflammatory response, and mares in the IUD-treatment group had the predicted number of occurrences of uterine oedema for the population size sampled. This, along with the serum progesterone and oestradiol data suggests that the IUD-treated mares were experiencing oestrous cycles.

Studies with ponies enabled more frequent observations of the IUDs and oestrous cycle events. From these observations we concluded that the 380 Copper “T” IUD was superior to the other IUDs tested for long-term contraception. In addition, observations on oestrous cycle events for one breeding season led us to conclude that cycle length for mares with IUDs was within the

normal range. The discrepancy in long-term rates of infertility between the mustang mares and pony mares equipped with a similar IUD is likely due to differences in uterine size. The retention of foreign objects in the mare uterus is related to the size of the object, relative to the size of the uterus. The ability of glass balls to be retained in the mare uterus has been shown to be related to the size of the glass ball (Nie et al. 2001; Thomas 2002). Although we have not found IUDs expelled by the uterus of an IUD-treated mare that became pregnant, we suspect the reason for the decline in efficacy of the Nevada mares was that the IUD was not retained in the uterus. This suggests that if a larger “T” IUDs was used, better rates of retention and contraceptive efficacy may be possible. Based on the pony studies, there is also evidence to suggest that shape of the IUD may also be a factor affecting retention, since neither the string GyneFix nor the ring IUDs were retained and performed as well as the T.

The mechanism preventing fertility in IUD-treated females has been argued to be either by interference with attachment of the early embryo to the uterus, or by induction of early abortion (Ortiz et al. 1996, Fourtney et al. 1999). This mechanism may differ among species and no data have been published for the mechanism in the mare. Because the oestrous cycle length of pony mares in this study was within in the normal range, we suggest that infertility in most instances was the result of the IUD interfering with events occurring between fertilization and early embryo attachment. However, in one pony mare that was observed to be 50-60 days pregnant, in a subsequent examination she was not pregnant and the IUD was visualized. This indicated that abortion had occurred, but the IUD was retained. This observation raises the possibility that if pregnancy occurs followed by abortion, IUD expulsion may also occur. It is also possible that if pregnancy occurs but abortion does not, the IUD could be expelled with the placenta at

parturition. This appears to have occurred with one Nevada mare who we observed to have the IUD when she was pregnant, and she went on to foal.

We believe that these studies provide evidence that long-term contraception of mare is possible with the SpayVacTM PZP vaccine. Further improvements to the formulation of GonConTM that are now being tested in white-tailed deer suggest that rates of contraception similar to SpayVacTM are achievable with GonaConTM. Development of larger IUDs that are better suited to the mustang mare may be possible. Regardless of the approach used, if a high rate of contraception is achievable for multiple years, population models suggest that contraception alone or used in conjunction with removal programs have the potential to stabilize and reduce population growth as well as reduce wild horse management costs (Garret et al. 1992, Cameron et al. 2001, Bartholow 2004, Ballou et al. this volume, Kirkpatrick and Turner this volume). These observations lead us to conclude that population management of wild horses by single application, multiyear contraceptives will likely be possible in the near future for horse populations that can be gathered from their range for treatment and release.

References

Ballou et al. this volume

Bartholow, J.M. (2004), An economic analysis of alternative fertility control and associated management techniques for three BLM wild horse herds: U.S. Geological Survey, Biological Resources Discipline, Open File Report 2004-119, 33 p. Brown, R. G., Bowen, W. D.,

Eddington, J. D., Kimmins, W. C., Mezei, M., Parsons, J. L. and B. Pohajdak (1997). Evidence for a long-lasting single administration vaccine in wild grey seals. *Journal of Reproductive Immunology* **35**, 43-51.

Burton, G.F., Szakal, A.K., Kapasi Z.F. and J. G. Tew. (1994). The generation and maintenance of antibody and B cell memory: The role of retained antigen and follicular dendritic cells. In *Strategies in Vaccine Design*, G.L. Ada (ed). Austin, Texas, R. G. Landes, pp 35-50.

Cameron, E.Z., Linklater, W.L., Minot, E.O. and K. J. Stafford (2001). Population dynamics 1994-98, and management of Kaimanawa wild horses. *Science for Conservation* **171**. viii + 165p.

Crowell-Davis, S. L. (2007) Sexual behavior of mares. *Hormones and Behavior* **52**, 12-17.

Curtis, P.D., Richmond, M.E., Miller, L.A., and F.W. Quimby (2007). Pathophysiology of white-tailed deer vaccinated with porcine zona pellucida immunocontraceptive. *Vaccine* **25**, 4623-4630.

Daels, P. F. and J. P. Hughes (1995). Fertility control using intrauterine devices: an alternative for population control in wild horses. *Theriogenology* **44**, 629-639.

Dalin, A.M., Anderson, O. and L. Malmgren (2002). Immunization against gnRH in mature

mares: antibody titers, ovarian function, hormonal levels and oestrous behaviour.

Journal of Veterinary Medicine and Applied Physiology, Pathology and Clinical Medicine **49**, 125-131.

Delves, P.J. and I.M. Roitt (2005). Vaccines for the control of reproduction status in mammals, and aspects of comparative interest. *Developmental Biology (Basel)* **121**, 265-273.

Elhay, M., Newbold, A. Britton, A., Turley, P., Dowsett, K and J. Walker (2007). Suppression of behavioural and physiological oestrus in the mare by vaccination against GnRH. *Australian Veterinary Journal* **85**, 39-45.

Fagerstone, K. A., Coffey, M.A., Curtis, P.D., Dolbeer, R.A., Killian, G.J., Miller, L.A. and L. M. Wilmot (2002). Wildlife Fertility Control. *Wildlife Society Technical Review* **02-2**, 29 pp.

Fagerstone et al. this volume

Fisher, C. (1983). The wild horse dilemma: background, prospective, and action. *Rangelands* **5**, 164-166.

Fortney, J.A., Feldblum, P.J. and E.J Raymond (1999). Intrauterine devices: The optimal long-term contraceptive method? *Journal of Reproductive Medicine* **44**, 269-273.

Fraker, M. A., Brown, R. G., Gaunt, G. E., Kerr, J. A. and B. Pohajdak (2002). Long-lasting,

single-dose immunocontraception of feral fallow deer in British Columbia. *Journal of Wildlife Management* **66**, 1141-1147.

Ginther, O.J. (1992) Reproductive Biology of the Mare; Basic and Applied Aspects. 2nd Edition. Equiservices Publishing, Cross Plains, Wisconsin, U.S.A. pp. 238-240.

Imboden, I., Janett, F., Burger, D., Crowe, M. A., Hassig, M. and R. Thun (2006). Influence of immunization against GnRH on reproductive cyclicity and estrous behavior in the mare. *Theriogenology* **66**, 1866-1875.

Killian, G., Miller, L., Diehl, N. K. Rhyan, J. and D.Thain (2004). Evaluation of three contraceptive approaches for population control of wild horses. *Proceedings of the Vertebrate Pest Conference* **21**, 263-268.

Killian, G., Eisemann, J., Wagner, D., Werner, J., Shaw, D., Engeman, R. and L. Miller (2006a). Safety and toxicity evaluation of GonaConTM immunocontraceptive vaccine in white-tailed deer (*Odocoileus virginianus*). *Proceedings of the Vertebrate Pest Conference* **22**, 82-87.

Killian, G., Miller, L., Diehl, N. K. Rhyan, J. and D. Thain (2006b). Long-term efficacy of three contraceptive approaches for population control of wild horses. *Proceedings of the Vertebrate Pest Conference* **22**, 67-71.

Killian, G., Miller, L., Rhyan, J. and H. Doten (2006c) Immunocontraception of Florida feral swine with a single-dose GnRH vaccine. *American Journal of Reproductive Immunology* **55**, 378-384.

Kirkpatrick, J. F., Liu, I. K. M. and J. W. Turner (1990). Remotely-delivered immunocontraception in feral horses. *Wildlife Society Bulletin* **18**, 326-330.

Kirkpatrick, J. F., Liu, I. K. M. Turner, J. W, Jr., Naugler, R. and R. Keiper (1992). Long-term effects of porcine zona pellucida immunocontraception on ovarian function of feral horses (*Equus caballus*). *Journal of Reproduction and Fertility* **94**, 437-444.

Kirkpatrick, J. F., R. Nuagle, I. K. M. Liu, J. W. Turner and M. Bernoco (1995). Effects of seven consecutive years of porcine zona pellucida contraception on ovarian function in feral mares. *Biology of Reproduction Monograph Series 1, Equine Reproduction* **VI**. 411-418.

Kirkpatrick J. F., Turner, J. W. Jr., Liu I. K., Fayrer-Hosken, R. and A. T. Rutberg (1997). Case studies in wildlife immunocontraception: wild and feral equids and white-tailed deer. *Reproduction, Fertility and Development* **9**, 105-110

Kirkpatrick, J. F. and A. Turner (2002). Reversibility of action and safety during pregnancy of immunization against porcine zona pellucida in wild mares (*Equus caballus*). *Reproduction Supplement*. **60**:197-202.

Kirkpatrick, J F. and A. Turner (2003). Absence of effects from immunocontraception on seasonal birth patterns and foal survival among barrier island wild horses. *Journal of Applied Welfare Science* **6**, 301-308.

Kirkpatrick and Turner, this volume

Levy, J., Miller, L. A., Cynda-Crawford, P., Ritchey, J. W., Ross, M. K. and K. A. Fagerstone (2004). GnRH immunocontraception of male cats. *Theriogenology* **62**, 1116-30.

Liu, I. K., Turner, J. W., Jr., van Leeuwen, E. M., Flanagan, D. R., Hedrick, J. L., Murata, K., Lane, V. M. and M. P. Morales-Levy (2005). Persistence of anti-zonae pellucidae antibodies following a single inoculation of porcine zonae pellucidae in domestic equine. *Reproduction* **129**, 181-190.

Lou, Y.H., Bagavant, H., Ang, J., McElveen, M.F., Thai, H.,and K. S. Tung (1996). Influence of autoimmune ovarian disease pathogenesis on ZP3 contraceptive vaccine design. *Journal of Reproduction and Fertility Supplement* **50**, 159-63.

Mahi-Brown, C. A., Yanagimachi, R., Nelson, M.L., Yanagimachi, H. and N. Palumbo (1988) Ovarian histopathology of bitches immunized with porcine zonae pelucida. *Journal of Reproductive Immunology and Microbiology* **18**, 94-103.

Massei et al. this volume

McCann, S. M., Marubayashi, U., Sun, H.Q. and W. H. Yu (1993). Control of follicle-stimulating hormone and luteinizing hormone release by hypothalamic peptides. *Annals New York Academy of Sciences* **687**, 55-59.

McCann, S.M, Kimura, M., Walczewska, A., Karanth, S., Rettori, S. and W. H. Yu (1998). Hypothalamic control of gonadotropin secretion by LHRH, FSHRF, NO, cytokines, and leptin. *Domestic Animal Endocrinology* **15**, 333-344.

McNeilly, A.S., Crawford, J.L., Taragnat, C., Nicol, L and J.R. McNeilly (2003). The differential secretion of FSH and LH: regulation through genes, feedback and packaging. *Reproduction, Supplement* **61**, 463-476.

Miller, L. A., Johns, B. E. and G. J. Killian. (1999). Long-term effects of PZP immunization on reproduction in white-tailed deer. *Vaccine* **18**: 568-574.

Miller, L. A., Johns, B. E. and G. J Killian (2000). Immunocontraception of white-tailed deer with GnRH vaccine. *American Journal of Reproductive Immunology* **44**, 266-274.

Miller, L., Crane K., Gaddis S. and G.J. Killian (2001). Porcine zona pellucida immunocontraception: Long-term health effects on white-tailed deer. *Journal of Wildlife Management* **65L**, 941-945.

Miller, L.A. and G. J. Killian (2001). Seven years of white-tailed deer immunocontraceptive research at Penn State University: A comparison of two vaccines. *Proceedings of the Wildlife*

Damage Management Conference **9**, 60-69.

Miller, L. A., Rhyan, J and G.J. Killian (2004). GonaCon™, a versatile GnRH contraceptive for a large variety of pest animal problems. *Proceedings of the Vertebrate Pest Conference* **21**, 269-273.

Naz, R K., Gupta, S. K., Gupta, J. C., Vyas, H. K. and G. P. Talwa (2005). Recent advances in contraceptive vaccine development: a mini-review. *Human Reproduction* **20**, 3271-3283.

Nie, G.J., Johnson, C.E. and J.G.W. Wenzel (2001). Use of a glass ball to suppress behavioral estrus in mares. *American Association of Equine Practitioners Proceedings* **47**, 246-248.

Ortiz, M.E., Croxatto H.B. and C.W. Bardin (1996). Mechanisms of action of intrauterine devices. *Obstetrics and Gynecology Survey Supplement* **12**, **51**, S42-S51.

Padmanabhan, V. and A.S. McNeilly (2001). Is there an FSH-releasing factor? *Reproduction* **121**, 21-30.

Perry, K. R., Ario, W. M., Bynum, K.S. and L. A. Miller (2006). GnRH single-injection immunocontraception of black-tailed deer. *Proceedings of the Vertebrate Pest Conference* **22**, 72-77.

Powell, D. M. (1999). Preliminary evaluation of porcine zona pellucida (PZP) immunocontraception for behavioral effects in feral horses (*Equus caballus*). *Journal of Applied Animal Welfare Science* **2**, 321-325.

Sample, J. (1997). Ultrasonic appearance and the pattern of uterine edema to time ovulation in mares. *American Association of Equine Practitioners Proceedings* **43**, 189-191.

Skinner, S. M., Mills, T., Kirchick, H. J. and B. S. Dunbar (1984). Immunization with zona pellucida proteins results in abnormal ovarian follicular differentiation and inhibition of gonadotropin-induced steroid secretion. *Endocrinology* **115**, 2418-32.

Stoops M. A., Liu, I. K., Shideler, S. E., Lasley, B. L., Fayrer-Hosken, R. A., Benirschke, K., Murata, K., van Leeuwen, E. M. and G. B. Anderson (2006). Effect of porcine zonae pellucidae immunisation on ovarian follicular development and endocrine function in domestic ewes (*Ovis aries*). *Reproduction, Fertility and Development* **18**, 667-76.

Thomas, H.S. (2002). Marbles for mares; An alternative for preventing estrus in mares. *American Quarter Horse Journal*. December 84-87.

Turner, J. W., Jr., Liu, I. K. and J. F. Kirkpatrick (1996). Remotely delivered immunocontraception in free-roaming feral burros (*Equus asinus*). *Journal of Reproduction and Fertility* **107**, 31-35.

Turner, J. W., Jr., I. K. M. Liu, D. R. Flanagan, A. T. Rutberg and J. F. Kirkpatrick (2000).

Immunocontraception in feral horses: a single-inoculation vaccine providing one year of infertility. *Journal of Wildlife Management* **65**, 235-241.

Turner, A. and J. F. Kirkpatrick (2002). Effects of immunocontraception on population longevity and body condition in wild mares (*Equus caballus*) *Reproduction Supplement* **60**: 187-195.

Turner, J.W., Jr., I.K. Liu, D.R. Flanagan, K.S. Bynum and A.T. Rutberg (2002). Porcine zona pellucida (PZP) immunocontraception of wild horses (*Equus caballus*) in Nevada: a ten year study. *Reproduction Supplement* **60**, 177-186.

Wu, S., Hu, J. and D. Wildemeersch (2000). Performance of the frameless GyneFix and the TCu380A IUDs in a 3-year multicenter, randomized, comparative trial in parous women. *Contraception* **61**, 91-98.

Yu, W. H, Karanth, S., Walczewska, A., Sower, S. A. and S. M.McCann (1997). A hypothalamic follicle-stimulating hormone-releasing decapeptide in the rat. *Proceedings of the National Academy of Sciences, USA*. **94**, 9499-9503.

Figure Legends

Figure 1. Comparative rates of contraception for the three methods of contraception for each year of the four year study. Rates of contraception for SpayVac treated mares for years one, two, three and four were 100% (12/12 mares), 83% (10/12), 83% (10/12) and 83% (10/12), respectively. Rates of contraception for GonaConTM treated mares were 93% (14 /15 mares), 64% (9/14), 57% (8/14) and 43% (6/14) for years one through four, respectively. Contraception rates for IUD-treated mares were 80% (12/15 mares), 29% (4/14), 14% (2/14) and 0% (0/14), respectively in years one through four of the study. In year two, one mare died in each of the GonaConTM- and IUD-treatment groups of causes not related to the treatments.

Figure 2. Average anti-PZP titers (\pm SEM) for the fall bleed for mares that were contracepted compared to titers of mares that became pregnant. The average titers for each of years 1-4 were for the serum samples for the 10 mares that were infertile. The titer for the pregnant mares was the average for the 2 mares that became pregnant during the study.

Figure 3. Average anti-GnRH titers (\pm SEM) for the fall bleed for mares that were contracepted compared to titers of mares that became pregnant. The sample sizes for titers for years 1-4 were 14, 9, 8 and 6, respectively. The average titer for the pregnant mares was for the 8 mares that became pregnant during the four year study.

Figure 4. Average progesterone values ($\mu\text{g/ml}$) (\pm SEM) in non-pregnant GonaConTM-treated mares (n = 35) and SpayVacTM-treated mares (n = 46) and non-pregnant IUD-treated mares (n = 17) for the fall bleed for all years of the study. Values for GonaConTM- and SpayVacTM-treated mares were significantly lower ($p < .02$) than values for IUD-treated mares.

Figure 5. Average serum progesterone ($\mu\text{g/ml}$) at the fall bleed for pregnant GonaConTM-treated- mares (n = 8), pregnant-SpayVacTM-treated mares (n = 2) and pregnant-IUD-treated mares (n = 7). Values were significantly different between IUD and GonaConTM-treated mares ($p < .025$); a comparison could not be run with the SpayVacTM because of only two observations.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5

Table 1. Percentage of reproductive tracts with edema revealed by ultrasonography of mares for each of the treatments compared to the predicted number of mares expected to be in estrus.

Percent	Year 1	Year 2	Year 3	Year 4
Predicted	25-30%	25-30%	25-30%	25-30%
IUD Actual	20%(2/10)	0%(0/4)	0%(0/2)	-
GonCon Actual	23% (3/13)	25% (2/8)	25% (2/8)	17%(1/6)
SpayVac Actual	82% (9/11)	91% (10/11)	100 % (10/10)	70%(7/10)

Table 2. Summary of IUD types installed in pony mares, dates of installation, duration of contraception and reason for failure.

IUD type/mare	Installation date	Last observation	Duration of contraception	Reason for failure
GyneFix				
Maddie	5/24/02	6/25/02	30d	Pregnant
Godiva	5/24/02	7/23/02	60d	Pregnant
Sprite	5/24/02	4/28/07	5 y	NA
Libby(2)	9/23/05	4/23/06	7 m	Pregnant
Ring				
Maddie	7/2/02	8/23/02	10d	Pregnant
Mandy	9/14/02	12/10/03	1 y+?	No data
Libby	5/25/04	7/01/04	36d	Pregnant
Libby	7/09/04	9/18/04	2 m	Pregnant
“T”				
Dewdrop	5/24/02	11/01/06	5 y	pyometria
Maddie	5/30/03	4/28/07	4 y	NA
Godiva	7/14/03	4/28/07	4 y	NA

Libby	5/24/02	5/15/03	1 y	Pregnant
Connie	5/25/04	4/28/07	3 y	NA
Wanda	9/04/02	5/30/03	1 y+?	No data
Remy	9/04/02	5/30/03	1 y+?	No data

Table 2

Table 3. Number of estrous cycles observed from April through August of 2004 in pony mares equipped with either the GyneFix or the 380 copper “T” IUD. All mares were mature except Connie, who was an 11 month old filly when she received the IUD in 2004.

Mare: IUD Type	Estrous Cycles
	April-August 2004
Sprite: GyneFix	6
Dewdrop: T	5
Maddie: T	4
Godiva: T	4
Connie: T	3
